arXiv:1410.8657v1 [math.AG] 31 Oct 2014

ON THE COHOMOLOGY OF STOVER SURFACE

AMIR DŽAMBIĆ, XAVIER ROULLEAU

Abstract. We study a surface discovered by Stover which is the surface with minimal Euler number and maximal automorphism group among smooth arithmetic ball quotient surfaces. We study the natural map $\wedge^2 H^1(S,\mathbb{C}) \to H^2(S,\mathbb{C})$ and we discuss the problem related to the so-called Lagrangian surfaces. We obtain that this surface S has maximal Picard number and has no higher genus fibrations. We compute that its Albanese variety A is isomorphic to $(\mathbb{C}/\mathbb{Z}[\alpha])^7$, for $\alpha = e^{2i\pi/3}$.

1. Introduction

By the recent work of M. Stover [14], the number of automorphisms of a smooth compact arithmetic ball quotient surface $X = \Gamma\backslash\mathbb{B}_2$ is bounded by $288 \cdot e(X)$, where $e(X)$ denotes the topological Euler number of X.
Furthermore, Stover characterizes the arithmetic ball quotient surfaces X whose automorphism groups attain this bound, which by analogy with Hurwitz curves, he calls *Hurwitz ball quotients*; all such surfaces are finite Galois coverings of the Deligne-Mostow orbifold $\Lambda\backslash\mathbb{B}_2$ corresponding to the quintuple $(2/12, 2/12, 2/12, 7/12, 11/12)$ (see [12, 14]).
Stover constructs also a Hurwitz ball quotient S with Euler number $e(S) = 63$ and automorphism group $\mathrm{Aut}(S)$ isomorphic to $U_3(3) \times \mathbb{Z}/3\mathbb{Z}$, of order $18144 = 2^5 3^4 7$. He shows that S is the unique Hurwitz ball quotient with Euler number $e = 63$, and moreover that $e = 63$ is the minimal possible value for the Euler number of a Hurwitz ball quotient. Having this property the surface S can be seen as the 2-dimensional analog of the Klein's quartic which is the unique curve uniformized by the ball $\mathbb{B}_1$ with minimal genus and maximal possible automorphism group.
Our aim is to study more closely the cohomology of this particular surface S, which we will call *Stover surface* in the following. This surface S has the following numerical invariants (see [14]):

$e(S)$	$H_1(S,\mathbb{Z})$	q	$p_g = h^{2,0}$	$h^{1,1}$	$b_2(S)$
63	$\mathbb{Z}^{14}$	7	27	35	89

Let V be a vector space. Let us recall that a 2-vector $w \in \wedge^2 V$ has *rank 1* or is *decomposable* if there are vectors $w_1, w_2 \in V$ with $w = w_1 \wedge w_2$. A vector $w \in \wedge^2 V$ has *rank* 2 if there exist linearly independent vectors $w_i \in V$, $i = 1, .., 4$ such that $w = w_1 \wedge w_2 + w_3 \wedge w_4$.

Let B be an Abelian fourfold and let $p : S \to B$ be a map such that $p(S)$ generates B. We say that S is *Lagrangian with respect to* p if there exists a basis $w_1, \ldots, w_4$ of $p^* H^0(B, \Omega_B)$ such that the rank 2 vector $w = w_1 \wedge w_2 + w_3 \wedge w_4$ is in the kernel of the natural map $\phi^{2,0} : \wedge^2 H^0(S, \Omega_S) \to H^0(S, K_S)$.

Theorem 1. *The surface S has maximal Picard number. The natural map*

$$\phi^{1,1} : H^0(S, \Omega_S) \otimes H^1(S, \mathcal{O}_S) \to H^1(S, \Omega_S)$$

is surjective with a 14-dimensional kernel. The kernel of the map

$$\phi^{2,0} : \wedge^2 H^0(S, \Omega_S) \to H^0(S, K_S)$$

is 7-dimensional and contains no decomposable elements. The set of rank 2 vectors in $Ker(\phi^{2,0})$ is a quadric hypersurface.
There exists an infinite number (up to isogeny) of maps $p : S \to B$ (where B is an Abelian fourfold) such that S is Lagrangian with respect to p.
The Albanese variety of S is isomorphic to $(\mathbb{C}/\mathbb{Z}[\alpha])^7$, for $\alpha = e^{2i\pi/3}$.

By the Castelnuovo - De Franchis Theorem, the fact that there are no decomposable elements in $\wedge^2 H^0(S, \Omega_S)$ means that S has no fibration $f : S \to C$ onto a curve of genus $g > 1$. Moreover Theorem 1 implies that S has the remarkable feature that both maps

$$\phi^{2,0} : \wedge^2 H^{1,0}(S) \to H^{2,0}(S)$$
$$\phi^{1,1} : H^{1,0}(S) \otimes H^{0,1}(S) \to H^{1,1}(S)$$

have a non-trivial kernel. With Schoen surfaces (see [10, Remark 2.6]), this is the second example of surfaces enjoying such properties. For more on this subject, see e.g. [1, 5, 2, 8, 9].

We obtain these results using Sullivan's theory on the second lower quotient of the fundamental group $\pi_1(S)$ of S (see [4]).

For the motivation and a historic account of surfaces with maximal Picard number we refer to [3].

Aknowledgements We are grateful to Marston Conder and Derek Holt for their help in the computations of Theorem 3.

2. The Second lower central quotient of the fundamental group of S

Let $\Pi := \pi_1(X)$ be the fundamental group of a manifold X. The group $H_1(X, \mathbb{Z})$ is the abelianization of Π: $H_1(X, \mathbb{Z}) = \Pi/\Delta$ where $\Delta := [\Pi, \Pi]$ is the derived subgroup of Π, that is, the subgroup generated by all elements $[h, g] = g^{-1}h^{-1}gh$, $h, g \in \Pi$.
The second group in the lower central series $[\Delta, \Pi]$ is the group generated by commutators $[h, g]$, with $h \in \Delta$, $g \in \Pi$. It is a normal subgroup of the commutator group Δ. According to [4], we have the following results:

Proposition 2. *(Sullivan) Let X be a compact connected Kähler manifold. There exists an exact sequence*

$$0 \to Hom(\Delta/[\Delta, \Pi], \mathbb{R}) \to \wedge^2 H^1(X, \mathbb{R}) \to H^2(X, \mathbb{R}).$$

(Beauville) Suppose $H_1(X, \mathbb{Z})$ is torsion free. Then the group $\Delta/[\Delta, \Pi]$ is canonically isomorphic to the cokernel of the map

$$\mu : H_2(X, \mathbb{Z}) \to Alt^2(H^1(X, \mathbb{Z})) \text{ given by } \mu(\sigma)(a, b) = \sigma \cap (a \wedge b),$$

where $Alt^2(H^1(X, \mathbb{Z}))$ is the group of skew-symmetric integral bilinear forms on $H^1(X, \mathbb{Z})$.

In the case of the Stover surface, computer calculations give us the following result:

Theorem 3. *Let $\Pi = \pi_1(S)$ be the fundamental group of the Stover surface and $\Delta = [\Pi, \Pi]$. The group $\Delta/[\Delta, \Pi]$ is isomorphic to $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}^{28}$.*

Proof. By the construction of S [14], the fundamental group Π is isomorphic to the kernel $ker(\varphi)$ of the unique epimorphism $\varphi : \Lambda \longrightarrow G$ from the Deligne-Mostow lattice Λ corresponding to the quintuple $(2/12, 2/12, 2/12, 7/12, 11/12)$ onto the finite group $G = U_3(3) \times \mathbb{Z}/3\mathbb{Z}$. The lattice Λ is described by Mostow in [12] as a complex reflection group, and by generators and relation by Cartwright and Steger in [7]. This lattice has presentation

$$\Lambda = \langle j, u, v, b | u^4, v^8, [u, j], [v, j], j^{-3}v^2, uvuv^{-1}uv^{-1}, (bj)^2(vu^2)^{-1}, [b, vu^2], b^3, (bvu^3)^3 \rangle.$$

MAGMA command `LowIndexSubgroups` is used to identify the unique subgroup $\Gamma \lhd \Lambda$ of index 3, which is $\Gamma = \langle u, jb, bj \rangle$. Using the primitive permutation representation of $U_3(3)$ of degree 28, MAGMA is able to identify an homomorphism φ from Γ onto $U_3(3)$ induced from the assignment

$$u \mapsto (3,8,23,20)(4,24,6,12)(7,9,14,22)(10,19,11,13)(15,16,21,18)(17,26,27,25)$$
$$jb \mapsto (1,9,20,12,19,23,6,16)(2,27,14,17,13,26,15,25)(3,24)(4,5,10,21,7,11,28,8)$$
$$bj \mapsto (1,13,20,15,19,2,6,14)(4,9,10,12,7,23,28,16)(5,27,21,17,11,26,8,25)(22,24).$$

This homomorphism extends to an homomorphism φ from Λ onto G such that $\Pi = ker(\varphi)$ is a torsion-free normal subgroup in Λ, it is the fundamental group of S (see [14]). Let be $\Delta = [\Pi, \Pi]$ and $\Delta_2 = [\Delta, \Pi]$. It is easy to check that that Δ_2 is distinguished into Π. The image of Δ under the quotient map $\Pi \longrightarrow \Pi/\Delta_2$ is Δ/Δ_2, but we observe that it is also equal the commutator subgroup $[\Pi/\Delta_2, \Pi/\Delta_2]$, and therefore, the computation of Δ/Δ_2 is reduced to the one of the derived group $[\Pi/\Delta_2, \Pi/\Delta_2]$.
The MAGMA command `g:=Rewrite(G,g)` is used to have generators and relations of both subgroups $\Gamma < \Lambda$ and $\Pi < \Gamma$. The command `NilpotentQuotient(.,2)` applied to Π describes Π/Δ_2 in terms of a polycyclic presentation. The derived subgroup $[\Pi/\Delta_2, \Pi/\Delta_2]$ is obtained with `DerivedGroup(.)` applied to Π/Δ_2. Finally, applying the MAGMA function `AQInvariants` to $[\Pi/\Delta_2, \Pi/\Delta_2]$, MAGMA computes that the structure of Δ/Δ_2 is $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}^{28}$. □

Corollary 4. *The dimension of the kernel of* $\wedge^2 H^1(S, \mathbb{R}) \to H^2(S, \mathbb{R})$ *is* 28.

3. Computation of the map $\wedge^2 H^1(S, \mathbb{C}) \to H^2(S, \mathbb{C})$

Let A be the Albanese variety of the Stover surface S. The invariants are:

$$H_1(A, \mathbb{Z}) = H_1(S, \mathbb{Z}) = \mathbb{Z}^{14},\ H_2(A, \mathbb{Z}) = \wedge^2 H_1(A, \mathbb{Z}),\ H^{2,0}(A) = \wedge^2 H^{1,0}(S)$$
$$H^{1,1}(A) = H^{1,0}(S) \otimes H^{0,1}(S),\ H^{0,2}(A) = \wedge^2 H^{0,1}(S),$$

and

$H_1(A, \mathbb{Z})$	q	$h^{2,0}(A)$	$h^{1,1}(A)$	$b_2(A)$
$\mathbb{Z}^{14}$	7	21	49	91

We have a map respecting Hodge decomposition

$$\begin{array}{ccccc} H^{2,0}(A) & \oplus & H^{1,1}(A) & \oplus & H^{0,2}(A) \\ \downarrow & & \downarrow & & \downarrow \\ H^{2,0}(S) & \oplus & H^{1,1}(S) & \oplus & H^{0,2}(S) \end{array},$$

which is an equivariant map of $\mathrm{Aut}(S)$-modules. By Corollary 4, the kernel of that map is 28 dimensional ; it is moreover a $\mathrm{Aut}(S)$-module.

According to the Atlas tables [11], the group $U_3(3)$ has 14 irreducible representations χ_i, $1 \leq i \leq 14$ of respective dimension $1, 6, 7, 7, 7, 14, 21, 21, 21, 27, 28, 28, 32, 32$.

The irreducible representations of $\mathrm{Aut}(S) = U_3(3) \times \mathbb{Z}/3\mathbb{Z}$ are the χ_i^t, $i = 1, ..., 14$, $t = 0, 1, 2$ where $(g, s) \in U_3(3) \times \mathbb{Z}/3\mathbb{Z}$ acts on the same space as χ_i with action $(g, s) \cdot v = \alpha^s g(v)$ with$\alpha = e^{2i\pi/3}$ a primitive third root of unity.

Theorem 5. *The image of S by the Albanese map $\vartheta : S \to A$ is 2-dimensional.*
The map $H^{1,1}(A) \to H^{1,1}(S)$ is surjective, with a 14 dimensional kernel isomorphic to χ_6^0 as an $\mathrm{Aut}(S)$-module. We have $H^1(S, \mathbb{Z}) = \chi_3^1 \oplus \chi_3^2$ and $H^{1,1}(S) = \chi_1^0 \oplus \chi_3^0 \oplus \chi_{10}^0$, as $\mathrm{Aut}(S)$-modules.
The kernel of the natural map $\wedge^2 H^0(S, \Omega_S) \to H^0(S, K_S)$ is 7-dimensional, isomorphic to χ_3^0 as a $\mathrm{Aut}(S)$-module.
The surface S has maximal Picard number.
The Albanese variety A of S is isomorphic to $(\mathbb{C}/\mathbb{Z}[\alpha])^7$, for $\alpha = e^{2i\pi/3}$.

Since A is CM, it follows that S is Albanese standard, that is, the class of its image inside its Albanese variety A sits in the subring of $H^*(A, \mathbb{Q})$ generated by the divisor classes. That contrasts with the above mentioned Schoen surfaces, see [10].

Proof. Suppose that the image of S in A is 1-dimensional. Then there exists a smooth curve C of genus 7 with a fibration $f : S \to C$ and the map $\wedge^2 H^0(S, \Omega_S) \to H^0(S, K_S)$ is the 0 map and the kernel of $\wedge^2 H^1(S, \mathbb{C}) \to H^2(S, \mathbb{C})$ is at least 42 dimensional, which is impossible. Thus the image of S by the Albanese map $\vartheta : S \to A$ is 2-dimensional.

According to the Atlas character table [11], the possibilities for the $U_3(3)$-module $H_1(S, \mathbb{Z}) = H_1(A, \mathbb{Z}) = \mathbb{Z}^{14}$ are:

$$\chi_3^{\oplus 2}, \mathcal{R}_{\mathbb{Z}}(\chi_4) = \mathcal{R}_{\mathbb{Z}}(\chi_5) = \chi_4 \oplus \chi_5, \chi_4^{\oplus 2}, \chi_5^{\oplus 2} \text{ or } \chi_6$$

where $\mathcal{R}_{\mathbb{Z}}(\chi_j)$ is the restriction to $\mathbb{Z}$ of the 7-dimensional complex representation χ_j defined over $\mathbb{Z}[i]$. It cannot be $\chi_4^{\oplus 2}$ nor $\chi_5^{\oplus 2}$ because these are not is not defined over $\mathbb{Z}$ (some traces of elements are in $\mathbb{Z}[i] \setminus \mathbb{Z}$). We cannot have $H^1(S, \mathbb{Z}) = \chi_6$ since χ_6 remains irreducible, but $H^1(S, \mathbb{Z}) \otimes \mathbb{C} = H^{1,0} \oplus H^{0,1}$ is a Hodge decomposition on which the representation of $U_3(3)$ splits.

By duality, the kernel of $H^{2,0}(A) \to H^{2,0}(S)$ has same dimension d as the kernel of $H^{0,2}(A) \to H^{0,2}(S)$. Let k be the dimension of the kernel of the $U_3(3)$-equivariant map $H^{1,1}(A) \to H^{1,1}(S)$. We have $28 = k + 2d$, moreover since $h^{1,1}(S) = 35$ and $h^{1,1}(A) = 49$, we get $28 \geq k \geq 14$.

Let us suppose that $H^1(S, \mathbb{Z}) = \chi_4 \oplus \chi_5$. Then the representation $H^{1,1}(A)$ equals to $\chi_4 \otimes \chi_5 = \chi_1 + \chi_7 + \chi_{10}$ (of dimension $1 + 21 + 27$). An Abelian variety on which a finite group G acts possesses a G-invariant polarization (for example $\sum_{g \in G} g^* L$, where L is some polarization). Therefore the one dimensional $\mathrm{Aut}(S)$-invariant space of $H^{1,1}(A)$ is generated by the class of an ample divisor and the natural map $\vartheta^* : H^{1,1}(A) \to H^{1,1}(S)$ is injective on that subspace. Therefore the map ϑ^* has a kernel of dimension $k = 21$, 27 or 48. This is impossible because $k + 2d$ equals 28.

Hence, we have $H^1(S, \mathbb{Z}) = \chi_3^{\oplus 2}$ and moreover

$$H^{2,0}(A) = \wedge^2 \chi_3 = \chi_3 \oplus \chi_6$$

(the dimensions are $21 = 7 + 14$) and

$$H^{1,1}(A) = \chi_3^{\otimes 2} = \chi_1 \oplus \chi_3 \oplus \chi_6 \oplus \chi_{10}$$

($49 = 1{+}7{+}14{+}27$). By checking the possibilities, we obtain $k = 14$, $H^{1,1}(S) = \chi_1 \oplus \chi_3 \oplus \chi_{10}$, and the map $H^{1,1}(A) \to H^{1,1}(S)$ is surjective. The kernel of the map $H^{2,0}(A) \to H^{2,0}(S)$ is isomorphic to χ_3, of dimension 7, the action of $U_3(3)$ on $H^{2,0}(S)$ is then $H^{2,0}(S) = \chi_6 \oplus \chi$, where χ is a 13 dimensional representation.

Let $\sigma \in Aut(S) = U_3(3) \times \mathbb{Z}/3\mathbb{Z}$ be the order 3 automorphism commuting with every other element. It corresponds to an element $\sigma' \in \Lambda$ normalizing Π in Λ and such that the group Π' generated by Π and σ' contains Π with index 3. Using MAGMA, one find that we can choose $\sigma' = j^4$, where j is the order 12 element described in the proof of Theorem 3.
The quotient surface S/σ of S by σ is equal to $\mathbb{B}_2/\Pi'$. The fundamental group of S' is Π'/Π'_{tors} where Π'_{tors} is the subgroup of Π' generated by torsion elements. Using MAGMA, one find that Π' has a set of 8 generators with 7 of them which are torsion elements. Using these elements, we readily compute that Π'/Π'_{tors} is trivial. Therefore the space of one-forms on S that are invariant by σ is 0. Using the symmetries of $U_3(3)$, one see that σ acts on the tangent space $H^0(S, \Omega_S)^*$ as the multiplication by α or α^2. After possible permutation of σ and σ^2, we can suppose it is α.

We see that the representation of $Aut(S)$ on $H_1(S, \mathbb{Z})$ is $\chi_3^1 \oplus \chi_3^2$. The lattice $H_1(S, \mathbb{Z}) \subset H^0(S, \Omega_S)^*$ is moreover a $\mathbb{Z}[\alpha]$-module. The ring $\mathbb{Z}[\alpha]$ is a principal ideal domain, therefore $H_1(S, \mathbb{Z}) = \mathbb{Z}[\alpha]^7$ (for the choice of a certain basis) and A is isomorphic to $(\mathbb{C}/\mathbb{Z}[\alpha])^7$.

Therefore A has maximal Picard number and all the classes of $H^{1,1}(A)$ are algebraic. These classes remain of course algebraic under the map $H^{1,1}(A) \to H^{1,1}(S)$, which is surjective. Thus S is a surface with maximal Picard number. □

4. Lagrangian surfaces and Stover surface

Let B be an Abelian fourfold and let $p : S \to B$ be a map such that $p(S)$ generates B. Let us recall that S is Lagrangian with respect to p if there exists a basis $w_1, \dots, w_4$ of $p^*H^0(B, \Omega_B)$ such that the rank 2 vector $w = w_1 \wedge w_2 + w_3 \wedge w_4$ is in the kernel of the natural map $\phi^{2,0} : \wedge^2 H^0(S, \Omega_S) \to H^0(S, K_S)$. Let us now prove

Theorem 6. *The 7 dimensional space $Ker(\phi^{2,0})$ contains no decomposable elements. The algebraic set of rank 2 vectors in $Ker(\phi^{2,0})$ is a quadric $\tilde{Q} \subset Ker(\phi^{2,0})$.*
There exists an infinite number (up to isogeny) of maps $p : S \to B$ where B is an Abelian fourfold such that S is Lagrangian with respect to p.
There exists an infinite number (up to isogeny) of maps $p : S \to B$ where B is an Abelian fourfold such that

$$\tilde{Q} \cap p^*H^0(B, \wedge^2\Omega_B) = \{0\},$$

and for some of them we even have $Ker(\phi^{2,0}) \cap p^*H^0(B, \wedge^2\Omega_B) = \{0\}$.
The generic rank 2 element w in $\tilde{Q} \subset Ker(\phi^{2,0})$ does not correspond to any morphism to an Abelian fourfold.

Proof. We proved in Theorem 5 that

$$H^{2,0}(A) = \wedge^2\chi_3 = \chi_3 \oplus \chi_6$$

and the kernel of $\phi^{2,0} : H^{2,0}(A) \to H^{2,0}(S)$ is the 7-dimensional subspace with representation χ_3. In a basis $\gamma = (e_1, \dots, e_7)$ of $\chi_3 = H^0(S, \Omega_S) = H^{1,0}(S)$, the two following matrices A, B are generators of the group $U_3(3)$:

$$A = \begin{pmatrix} -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}, B = \begin{pmatrix} 0 & -1 & 0 & 0 & 0 & -1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 1 \end{pmatrix}.$$

Using the basis $\beta = (e_{ij})_{1 \le i < j \le 7}$ of $\wedge^2\chi_3$ ($e_{ij} = e_i \wedge e_j$) with order $e_{ij} \le e_{st}$ if $i < s$ or $i = s$ and $j \le t$, one computes that the subspace $Ker(\phi^{2,0}) = \chi_3 \subset \wedge^2\chi_3$ is generated by the columns of the matrix $M \in M_{21,7}$, where ${}^tM = (N, 2I_7)$, for

$$N = \begin{pmatrix} 0 & 0 & 2 & -2 & -2 & -2 & 0 & 2 & 2 & -2 & 2 & 2 & 2 & 2 \\ -1 & 0 & 0 & 2 & 4 & 0 & 1 & -3 & -3 & 1 & -3 & -4 & -2 & -4 \\ 0 & -2 & 0 & -2 & -2 & 0 & -2 & 2 & 2 & 0 & 0 & 2 & 2 & 2 \\ -1 & -2 & 2 & 0 & -2 & 0 & -1 & 1 & 3 & 1 & 1 & 0 & 0 & 2 \\ -1 & 1 & -1 & 3 & 1 & 3 & 0 & -4 & -2 & 2 & 0 & -4 & -2 & -2 \\ 0 & 3 & -3 & 1 & -1 & 1 & 1 & -3 & -3 & -1 & 1 & 0 & -2 & -2 \\ 1 & 1 & 1 & 3 & 3 & 1 & 2 & -2 & 0 & 0 & 0 & -2 & 0 & -2 \end{pmatrix} \in M_{7,14}$$

and I_7 the 7×7 identity matrix. Knowing that, we obtain the ideal I_V of the algebraic set V of couples $(w_1, w_2) \in \chi_3 \oplus \chi_3$ such that $w_1 \wedge w_2 \in Ker(\phi^{2,0}) \subset \wedge^2\chi_3$. That ideal is generated by 14 homogeneous quadratic polynomials in the variables $x_1, \dots, x_{14}$. Let W be

the algebraic set of couples $(w_1, w_2) \in \chi_3 \oplus \chi_3$ such that $w_1 \wedge w_2 = 0 \in \wedge^2\chi_3$. The ideal I_W of W is generated by the 2 by 2 minors of the matrix

$$L = \begin{pmatrix} x_1 & \dots & x_7 \\ x_8 & \dots & x_{14} \end{pmatrix}.$$

Since $W \subset V$, we have $Rad(I_V) \subset Rad(I_W)$ where $Rad(I)$ is the radical of an ideal I. On the other hand, using Maple, one can check that the 21 minors of L are in $Rad(I_V)$, hence $Rad(I_W) \subset Rad(I_V)$, thus $V = W$.
We therefore conclude that the kernel of $\phi^{2,0}$ contains no decomposable elements.

A 2-vector w over a characteristic 0 field can be expressed uniquely as $w = \sum_{i,j} a_{ij} e_i \wedge e_j$ where $a_{ij} = -a_{ji}$. The rank of the vector w is half the rank of the (skew-symmetric) coefficient matrix $A_w := (a_{ij})_{1 \le i,j \le 7}$ of w [6, Thm 1.7 & Remark p. 13]. Thus the 2-vector $w = a_1 v_1 + \dots + a_7 v_7$ in $Ker(\phi^{2,0})$ (where the v_i, $i = 1..7$ are the vectors corresponding to the columns of the matrix M) is a rank 2 vector if and only if the 49 6×6 minors of the matrix A_w are 0. The radical of the ideal generated by these minors is principal, generated by a homogeneous quadric in $a_1, \dots, a_7$ whose associated symmetric matrix is

$$Q = \begin{pmatrix} 7 & 3 & 3 & 1 & -3 & -3 & -5 \\ 3 & 7 & 3 & 3 & 1 & -3 & -3 \\ 3 & 3 & 7 & 3 & 3 & 1 & -3 \\ 1 & 3 & 3 & 7 & 3 & 3 & 1 \\ -3 & 1 & 3 & 3 & 7 & 3 & 3 \\ -3 & -3 & 1 & 3 & 3 & 7 & 3 \\ -5 & -3 & -3 & 1 & 3 & 3 & 7 \end{pmatrix}.$$

Therefore $w \in Ker(\phi^{2,0})$ has rank 2 if and only if $(a_1, \dots, a_7) Q^t (a_1, \dots, a_7) = 0$.

The point $(10 + 8\alpha, -7, 0, 0, 7, 0, 0)$ lies on the associated smooth quadric $\tilde{Q}$, therefore $\tilde{Q}(\mathbb{Q}[\alpha])$ is infinite. Let be w be a 2-vector in $\tilde{Q}(\mathbb{Q}[\alpha])$. The decomposable vector $\wedge^2 w \neq 0$ has coordinates in $\mathbb{Q}[\alpha]$ in the basis $(e_{i1} \wedge . \cdots \wedge e_{i4})$ of $\wedge^4 H^0(S, \Omega_S)$. The corresponding 4-dimensional vector space W is therefore generated by 4 vectors $w_1, \dots, w_4$ with coordinates over $\mathbb{Q}[\alpha]$ in the basis $\gamma = (e_1, \dots, e_7)$ of $H^0(S, \Omega_S)$.
One computes that the image of $\mathbb{Q}[\alpha][U_3(3) \times \mathbb{Z}/3\mathbb{Z}]$ in $M_7(\mathbb{Q}[\alpha])$ is 49 dimensional over $\mathbb{Q}[\alpha]$, thus

$$\mathbb{Q}[U_3(3) \times \mathbb{Z}/3\mathbb{Z}] = M_7(\mathbb{Q}(\alpha)) \, (= End(A) \otimes \mathbb{Q})$$

in the basis γ, ($H_1(S, \mathbb{Q}[\alpha])$ is the $\mathbb{Q}[\alpha]$-vector space generated by $e_1, \dots, e_k$) and therefore there exists a morphism $p : S \to E^4 = B$ (where $E = \mathbb{C}/\mathbb{Z}[\alpha]$) such that $W = p^* H^0(B, \Omega_B)$. By hypothesis the image $p(S)$ generates B. By construction

$$\wedge^2 p^* H^0(B, \Omega_B) \cap Ker(\phi^{2,0})$$

is at least one dimensional since it contains w, and therefore S is Lagrangian for p.

A contrario, the trace of an order 2 automorphism $\sigma \in \mathrm{Aut}(S) \subset \mathrm{Aut}(A)$ acting on the tangent space of A at 0 equals to -1, therefore the image B' of the endomorphism $p : \sigma - 1_A$, where 1_A is the identity of A is an Abelian fourfold. Using Maple, one computes that

$$\wedge^2 p^* H^0(B, \Omega_B) \cap Ker(f) = \{0\}.$$

Let $\vartheta : S \to A$ be the Albanese map of S, and let $q : A \to A$ be an endomorphism with a 4 dimensional image and a representation in $M_7(\mathbb{Q}) \subset M_7(\mathbb{Q}(\alpha))$ in the basis γ. Since the matrix Q is positive definite, we have

$$\wedge^2 p^* H^0(B, \Omega_B) \cap \tilde{Q} = \{0\},$$

where p is the map $p = q \circ \vartheta : S \to B$. Therefore S is not Lagrangian with respect to p. □

Remark 7. Let X be a surface and let $\phi^{2,0} : \wedge^2 H^0(X, \Omega_X) \to H^0(X, K_X)$ be the natural map. Let be $d = \dim Ker(\phi^{2,0})$ and $q = \dim H^0(X, \Omega_X)$. In the proof of Theorem 6, we saw that the set of rank k vectors in $Ker(\phi^{2,0})$ is a determinantal variety: the intersection of minors of size $\geq 2k+1$ of some anti-symmetric matrix of size $q \times q$ with linear entries in d variables. It seems to the authors quite remarkable that for Stover's surface the set of rank 2 vectors (obtained as the zero set of 49 6×6 minors of a size $q = 7$ matrix) is an hypersurface in $Ker(\phi^{2,0})$. That hypersurface is the only $U_3(3)$-invariant quadric of $U_3(3)$ acting on $Ker(\phi^{2,0})$.

References

[1] Barja M.A., Naranjo J.C., Pirola G.P., *On the topological index of irregular surfaces*, J. Algebraic Geom. 16 (2007), no. 3, 435–458.

[2] Bastianelli F., Pirola G.P., Stoppino L., *Galois closure and Lagrangian varieties*, Advances in Math., 225 (2010), 3463–3501.

[3] Beauville A., *Some surfaces with maximal Picard number*, Journal de l'École Polytechnique 1 (2014), 101–116.

[4] Beauville A., *On the second lower quotient of the fundamental group of a compact Kahler manifold*, ArXiv, to appear in "Algebraic and Complex Geometry" in honor of K. Hulek (Springer).

[5] Bogomolov F., Tschinkel Y., *Lagrangian subvarieties of abelian fourfolds*, Asian J. Math. 4 (2000), no. 1, 19–36.

[6] Bryant R.L., Chern S.S., Gardner R.B., Goldschmidt H.H., Griffiths P.A. *Exterior differential systems*, (Mathematical Sciences Research Institute publications, 18), Springer-Verlag (1991).

[7] Cartwright D., Steger T., *Enumeration of the 50 fake projective planes*, Comptes Rendus Mathematique 348 (1): 11–13, doi:10.1016/j.crma.2009.11.016

[8] Campana, F. *Remarques sur les groupes de Kähler nilpotents*, Ann. Sci. École Norm. Sup. (4) 28 (1995), no. 3, 307–316.

[9] Causin A, Pirola G.P., *Hermitian matrices and cohomology of Kähler varieties*, Manuscripta Math. 121 (2006), no. 2, 157–168.

[10] Ciliberto C., Mendes Lopes M., Roulleau X., *On Schoen surfaces*, to appear in Comment. Math. Helvet.

[11] Conway, J., Curtis R., Parker R., Norton S., Wilson R., *Atlas of finite groups*. Oxford University Press, 1985.

[12] Mostow G.D., *On a remarkable class of polyhedra in complex hyperbolic space*, Pacific J. Math., 86(1) 1980, 171–276.

[13] Schoen C., *Albanese standard and Albanese Exotic varieties*, J. of London Math. Soc. (2) 74 (2006) 304–320.

[14] Stover M., *Hurwitz ball quotients*, Math. Z. 277 (2014), no. 1–2, 75–91

Amir Džambić,
Johann Wolfgang Goethe Universität, Institut für Mathematik,
Robert-Mayer-Str. 6-8, 60325 Frankfurt am Main,
Germany
dzambic@math.uni-frankfurt.de

Xavier Roulleau,
Laboratoire de Mathématiques et Applications, Université de Poitiers,
Téléport 2 - BP 30179 - 86962 Futuroscope Chasseneuil
France
roulleau@math.univ-poitiers.fr